

## **Medical Errors**

### **What are Medical Errors?**

A medical error is an inaccurate or incomplete diagnosis and/or treatment of a disease, injury, syndrome, behavior, infection or other ailment that causes serious harm to the patient.

Medical errors can occur anywhere in the health care system: Hospitals, clinics, outpatient surgery centers, doctors' offices, nursing homes, pharmacies, or patients' homes. These errors can involve medicines, surgery, diagnosis, equipment and lab reports. When an injury is caused by poor medical care rather than by the underlying disease or condition of the patient, it is known as an “adverse event”.

### **What are the consequences of medical errors?**

Medical errors are one of the nation's leading causes of death and injury. A report from the Institute of Medicine estimates that medical errors cause 44,000 to 98,000 unnecessary deaths and 1,000,000 excess injuries each year in the US<sup>1</sup>. It is estimated that 2.9% to 3.7% of hospital admissions involve an adverse event.<sup>2</sup> More people die from medical errors each year than from motor vehicle accidents, breast cancer, or AIDS.<sup>1</sup>

Medical errors also carry a high financial cost. It is estimated that medical errors cost the nation \$37.6 billion each year.<sup>1</sup>

The Connecticut Department of Public Health recorded 1,224 adverse events in CT hospitals in 2009. The most common errors were (1) falls resulting in serious disability or death- (41%) (2) Errors during endoscopy and laparoscopy (20%) (3) Pressure ulcers acquired after admission to a healthcare facility and (4) leaving objects in patients after surgery.<sup>3</sup>

### **What causes medical errors?**

Medical errors are more common with inexperienced physicians, new procedures, extremes of patient age, complex care and urgent care.<sup>4</sup> Poor communication, improper documentation, illegible handwriting, inadequate nurse-to-patient ratios, and similarly named medications are also contribute to the problem. Patients may also contribute significantly to medical errors. Falls, for example, are often due to patients' own misjudgments.

Sleep deprivation has also been cited as a contributing factor in medical errors. One study found that being awake for over 24 hours caused medical interns to double or triple the number of preventable medical errors, including those which resulted in injury or death<sup>5</sup>.

### **How can medical errors be prevented?**

Government agencies, purchasers of care, and providers must work together to make the health care system safer for patients and the public. Research shows that the majority of medical errors can be prevented. A study found that 70 percent of adverse events are preventable.<sup>6</sup>

Possible solutions include:

- 1) Use of information technology, such as hand-held bedside computers, to eliminate reliance on handwriting for ordering medications and other treatment needs. Computerized prescribing systems lower the probability that a physician will make errors by providing accurate information on the drug, on the patient and on possible drug interactions and allergy reactions.<sup>7</sup>
- 2) Avoidance of similar-sounding and look-alike names and packages of medication.
- 3) Standardization of treatment policies and protocols to avoid confusion and reliance on memory, which is known to be fallible and responsible for many errors.
- 4) Following simple checklists to improve health care safety.<sup>8</sup>

Brandon Levan  
CT Health Policy Project Senior Fellow, Board member  
May 2010

**Resources:**

- [1] Institute of Medicine. "To Err is Human: Building A Safer Health System." November 1,1999
- [2] Brennan TA, Leape LL, Laird NM, Hebert L, Localio AR, Lawthers AG, et al. Incidence of adverse events and negligence in hospitalized patients. Results of the Harvard Medical Practice Study I. *N Engl J Med* 1991;324(6):370-7.
- [3] The Connecticut Department of Public Health. Health Care Quality Reports. June 2009
- [4] Weingart, Saul; Ciara Wilson, Bob Marley, Karina (2000). "Epidemiology of medical error". BMJ Publishing Group. <http://bmj.bmjournals.com/cgi/content/full/320/7237/774>. Retrieved 2006-03-17.
- [5] Barger LK, Ayas NT, Cade BE, Cronin JW, Rosner B, et al. (2006) Impact of Extended-Duration Shifts on Medical Errors, Adverse Events, and Attentional Failures. *PLoS Med* 3(12): e487  
doi:10.1371/journal.pmed.0030487
- [6] Agency for Healthcare Research and Quality. "Medical Errors: The Scope of the Problem". <http://www.ahrq.gov/qual/errback.htm>
- [7] Nguyen, Giang. "Preventing Medical Errors: Use of Information Technology to Reduce Adverse Drug Events". Bright Hub. Dec 2008.

[8] Morgera, Vicent. "Checklist Can Reduce Fatal Medical Errors". 24-7 Press Release. March 05, 2010.  
<http://www.24-7pressrelease.com/press-release/checklists-can-reduce-fatal-medical-errors-139993.php>